

04/08/99



JC645 U.S. PTO

JC625 U.S. PTO

09/288344

04/08/99

UTILITY PATENT APPLICATION TRANSMITTAL
 (only for new and continuation-in-part
 nonprovisional applications under 37 CFR
 1.53(b))

Docket No.:

P-PM 3474

Address to: ASSISTANT COMMISSIONER FOR PATENTS
Box Patent Application
 Washington, D.C. 20231

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING LABEL NUMBER: EL 041 092 647 US

DATE OF DEPOSIT: April 8, 1999

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING
 DEPOSITED WITH THE UNITED STATES POSTAL SERVICE
 "EXPRESS MAIL POST OFFICE TO ADDRESSEE" SERVICE UNDER
 37 C.F.R. 1.10 ON THE DATE INDICATED ABOVE, AND IS
 ADDRESSED TO: **BOX PATENT APPLICATION, ASSISTANT**
COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231.

Rebecca McElroy

(FAXED OR PRINTED NAME OR PERSON MAILING PAPER OR FEE)

(SIGNATURE OF PERSON MAILING PAPER OR FEE)

This is a request for filing a

X new utility patent application under 37 CFR 1.53(b)(1).

_____ continuation-in-part under CFR 1.53(b)(2) of prior
 application serial no. _____, filed _____
 (list entire parentage).

Title: **METHODS OF OPTIMIZING DRUG THERAPEUTIC EFFICACY FOR
 TREATMENT OF IMMUNE-MEDIATED GASTROINTESTINAL DISORDERS**

Inventor(s) (full name of each inventor): **Ernest G. Seidman,
 Yves Théorêt**

Enclosed are:

- X Return receipt postcard
- X Initial Information Data Sheet
- X 1 Page application cover sheet
- X 50 Pages of specification (includes claims and abstract)
- X 3 Sheets of drawing(s)
- _____ Pages of an executed Declaration for Patent Application
- _____ An executed Power of Attorney for Patent Application by Assignee
- _____ Paper copy of sequence listing, pages _____ through _____
- _____ Sequence listing in computer readable form
- _____ Statement Under 37 CFR 1.821(f)
- _____ An executed assignment and cover sheet
- _____ An executed small entity statement
- _____ Also enclosed: _____

Inventors: **Seidman and Théorêt**
Docket No.: **P-PM 3474**
Page 2

— This application is based on prior foreign application(s) No. (s) _____, filed in _____ on _____, respectively, and priority is hereby claimed therefrom.

☒ This application is based on, and claims the benefit of, U.S. Provisional Application No. 60/101,714, filed September 24, 1998, and entitled METHODS OF OPTIMIZING DRUG THERAPEUTIC EFFICACY FOR TREATMENT OF INFLAMMATORY BOWEL DISEASE.

— This application is based on, and claims the benefit of, U.S. Provisional Application No. 60/_____ (yet to be assigned), filed _____, which was converted from U.S. Serial No. _____, and entitled _____.

The filing fee has been calculated as shown below:

	Number Filed		Number Extra		Rate			Fee	
					Small Entity	Other Entity		Small Entity	Other Entity
Total Claims	34-20	=	14	x	\$9	\$18	=	\$0	\$
Independent Claims	4-3	=	1	x	\$39	\$78	=	\$0	\$
Multiple Dependent Claims Presented: ___ Yes <input checked="" type="checkbox"/> No					\$130	\$260		\$0	\$
					BASIC FEE			\$380	\$760
					TOTAL FEE			\$0	\$

— A check in the amount of \$ _____ to cover the filing fee is enclosed.

☒ The payment of the filing fee is to be deferred until the Declaration is filed. Do not charge our deposit account.

— The Commissioner is hereby authorized to charge fees under 37 CFR 1.16 and 1.17 which may be required or credit any overpayment to Deposit Account No. _____. A duplicate copy of this sheet is enclosed.


Inventors: **Seidman and Théorêt**
Docket No.: **P-PM 3474**
Page 3

Address all future communications to:

Cathryn Campbell
CAMPBELL & FLORES LLP
4370 La Jolla Village Drive, Suite 700
San Diego, California 92122
telephone: (619) 535-9001
facsimile: (619) 535-8949

Respectfully submitted,

Date: April 8, 1999


Deborah L. Cadena
Registration No. 44,048
CAMPBELL & FLORES LLP
4370 La Jolla Village Drive
Suite 700
San Diego, California 92122

Initial Information Data Sheet

Inventor Information

Inventor One Given Name :: Ernest
 Family Name :: Seidman
 Name Suffix :: G.
 Postal Address Line One :: 5617 Hartwell
 Postal Address Line Two ::
 City :: Côte St. Luc
 State/Province :: Quebec
 Country :: Canada
 Postal or Zip Code :: H4W 1T5
 City of Residence :: Côte St. Luc
 State/Prov. of Residence :: Quebec
 Country of Residence :: Canada
 Citizenship :: Canadian

Inventor Two Given Name :: Yves
 Family Name :: Théorêt
 Name Suffix ::
 Postal Address Line One :: 4824 Harvard Avenue
 Postal Address Line Two ::
 City :: Montreal
 State/Province :: Quebec
 Country :: Canada
 Postal or Zip Code :: H3X 3N9
 City of Residence :: Montreal
 State/Prov. of Residence :: Quebec
 Country of Residence :: Canada
 Citizenship :: Canadian

Correspondence Information

Name Line One :: CAMPBELL & FLORES LLP
 Address Line One :: 4370 La Jolla Village Drive
 Address Line Two :: Suite 700
 City :: San Diego
 State/Province :: California
 Country :: USA
 Postal or Zip Code :: 92122
 Telephone :: (619) 535-9001
 Fax :: (619) 535-8949
 E-Mail :: CColacino@candf.com

Application Information

Title Line One :: METHODS OF OPTIMIZING DRUG THERAPEUTIC
 Title Line Two :: EFFICACY FOR TREATMENT OF
 Title Line Three :: IMMUNE-MEDIATED GASTROINTESTINAL
 Title Line Four :: DISORDERS
 Total Drawing Sheets :: 3
 Formal Drawings :: No
 Application Type :: Utility
 Docket Number :: P-PM 3474

Representative Information

Registration Number One :: 31,815
 Registration Number Two :: 36,405
 Registration Number Three :: 34,949
 Registration Number Four :: 30,806
 Registration Number Five :: 38,701
 Registration Number Six :: 36,933
 Registration Number Seven :: 39,200
 Registration Number Eight :: 38,444
 Registration Number Nine :: 37,915
 Registration Number Ten :: 41,029
 Registration Number Eleven :: 44,048

Continuity Information

This application is a :: New Application which Claims Benefit of
 > Application One :: 60/101,714
 Filing Date :: September 24, 1998
 which is a :: Provisional Application

 >> Application Two :: [ENTER SERIAL NUMBER]
 Filing Date :: [ENTER FILING DATE]
 Patent Number :: [ENTER PATENT NO OR DELETE ENTIRE LINE]
 which is a :: [ENTER Continuation, Continuation in
 Part, File Wrapper Continuation,
 Division] of

>>> Application Three :: [ENTER SERIAL NUMBER]
Filing Date :: [ENTER FILING DATE]
Patent Number :: [ENTER PATENT NO OR DELETE ENTIRE LINE]

Prior Foreign Applications

Foreign Application One :: [ENTER APPLICATION NO.]
Filing Date :: [ENTER DATE]
Country :: [ENTER COUNTRY]
Priority Claimed :: [ENTER Y OR N]

0
1
2
3
4
5
6
7
8
9
A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y
Z

A P P L I C A T I O N

for

UNITED STATES LETTERS PATENT

on

METHODS OF OPTIMIZING DRUG THERAPEUTIC EFFICACY FOR
TREATMENT OF IMMUNE-MEDIATED GASTROINTESTINAL DISORDERS

by

Ernest G. Seidman

and

Yves Théorêt

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING LABEL NUMBER: EL 041 062 647 US

DATE OF DEPOSIT: April 8, 1999

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING
DEPOSITED WITH THE UNITED STATES POSTAL SERVICE
"EXPRESS MAIL POST OFFICE TO ADDRESSEE" SERVICE UNDER
37 C.F.R. 1.10 ON THE DATE INDICATED ABOVE, AND IS
ADDRESSED TO: BOX PATENT APPLICATION, ASSISTANT
COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231

Rebecca McElroy
(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)

(SIGNATURE OF PERSON MAILING PAPER OR FEE)

Sheets of Drawings: Three

Docket No.: P-PM 3474

Attorneys

CAMPBELL & FLORES LLP
4370 La Jolla Village Drive, Suite 700
San Diego, California 92122

**METHODS OF OPTIMIZING DRUG THERAPEUTIC EFFICACY FOR
TREATMENT OF IMMUNE-MEDIATED GASTROINTESTINAL DISORDERS**

This application claims the benefit of priority
of provisional application serial No. 60/101,714, filed
5 September 24, 1998, which is incorporated herein by
reference.

BACKGROUND OF THE INVENTION

The present invention relates generally to
autoimmunity and immune-mediated gastrointestinal
10 disorders such as inflammatory bowel disease and more
specifically to methods for optimizing treatment of
immune-mediated gastrointestinal disorders.

BACKGROUND INFORMATION

Immune-mediated gastrointestinal disorders
15 encompass a wide range of debilitating gastrointestinal
diseases of various etiologies. One such immune-mediated
gastrointestinal disorder, inflammatory bowel disease
(IBD), is the collective term used to describe two
gastrointestinal disorders of unknown etiology: Crohn's
20 disease (CD) and ulcerative colitis (UC). The course and
prognosis of IBD, which occurs world-wide and is reported
to afflict as many as two million people, varies widely.
Onset of IBD is predominantly in young adulthood with
diarrhea, abdominal pain, and fever the three most common
25 presenting symptoms. The diarrhea may range from mild to
severe and in ulcerative colitis often is accompanied by
bleeding. Anemia and weight loss are additional common
signs of IBD. Ten percent to fifteen percent of all
patients with IBD will require surgery over a ten year
30 period. In addition, patients with IBD are at increased

risk for the development of intestinal cancer. Reports of an increasing occurrence of psychological problems, including anxiety and depression, are perhaps not surprising symptoms of what is often a debilitating disease that strikes people in the prime of life.

6-Mercaptopurine (6-MP) and azathioprine (AZA), a pro-drug that is non-enzymatically converted to 6-mercaptopurine (6-MP), are 6-MP drugs that can be used as an effective treatment for inflammatory bowel diseases such as Crohn's disease and ulcerative colitis (Kirschner Gastroenterology 115:813-821 (1998)). 6-MP can be enzymatically converted to various 6-MP metabolites, including 6-methyl-mercaptopurine (6-MMP) and 6-thioguanine (6-TG) and their nucleotides. 6-TG nucleotides are thought to be the active metabolite in mediating many of the effects of 6-MP drug treatment.

Thiopurine methyltransferase (TPMT) is a cytoplasmic enzyme that preferentially catalyzes the S-methylation of 6-MP and 6-TG to form S-methylated metabolites such as 6-MMP and 6-methylthioguanine (6-MTG), respectively. TPMT exhibits genetic polymorphism, with 89% of Caucasians and African Americans having high activity, 11% intermediate activity and 1 in 300 TPMT deficient. Clinical studies with AZA and 6-MP have shown an inverse relationship between TPMT activity and 6-TGN accumulation. Patients who less efficiently methylate these thiopurines have more extensive conversion to 6-TGN, which can lead to potentially fatal hematopoietic toxicity. Therefore, patients who have less active TPMT can be more susceptible to toxic side effects of 6-MP therapy.

Although drugs such as 6-MP and AZA have been used for treating IBD, non-responsiveness and drug toxicity unfortunately complicate treatment in some patients. Complications associated with 6-MP drug treatment include allergic reactions, neoplasia, opportunistic infections, hepatitis, bone marrow suppression, and pancreatitis. Therefore, many physicians are reluctant to treat patients with AZA because of its potential side effects, especially infection and neoplasia.

Thus, there exists a need to develop methods to optimize the dose of 6-mercaptopurine drugs and assess biotransformation in individual patients to optimize the therapeutic efficacy of 6-mercaptopurine drugs while minimizing toxic side effects. The present invention satisfies this need and provides related advantages as well.

SUMMARY OF THE INVENTION

The present invention provides a method of optimizing therapeutic efficacy of 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder. The method includes the steps of administering a 6-mercaptopurine drug to a subject having an immune-mediated gastrointestinal disorder; and determining a level of 6-thioguanine in the subject having the immune-mediated gastrointestinal disorder, where a level of 6-thioguanine less than a level corresponding to about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of 6-mercaptopurine drug subsequently administered to the subject and where a level of 6-thioguanine greater than a level corresponding to about 400 pmol per 8×10^8 red blood cells indicates a need to

decrease the amount of 6-mercaptopurine drug subsequently administered to the subject. The methods are directed to treating immune-mediated gastrointestinal disorders, including inflammatory bowel diseases (IBD) such as

5 Crohn's disease and ulcerative colitis, lymphocytic colitis, microscopic colitis, collagenous colitis, autoimmune enteropathy, allergic gastrointestinal disease and eosinophilic gastrointestinal disease. In a method of optimizing therapeutic efficacy of 6-mercaptopurine

10 treatment of IBD, the subject having IBD can be, for example, a pediatric subject. The level of 6-thioguanine can be determined, for example, in red blood cells using high pressure liquid chromatography.

The present invention also provides a method of

15 reducing toxicity associated with 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder. The method of reducing toxicity associated with an immune-mediated gastrointestinal disorder includes the steps of administering a 6-mercaptopurine

20 drug to a subject having the immune-mediated gastrointestinal disorder; and determining a level of a 6-mercaptopurine metabolite in the subject having the immune-mediated gastrointestinal disorder, where a level of the 6-mercaptopurine metabolite greater than a

25 predetermined toxic level of the 6-mercaptopurine metabolite indicates a need to decrease the amount of 6-mercaptopurine drug subsequently administered to the subject, thereby reducing toxicity associated with 6-mercaptopurine drug treatment of the immune-mediated

30 gastrointestinal disorder. In a method of the invention, the 6-mercaptopurine metabolite can be, for example, 6-thioguanine and the predetermined toxic level of 6-thioguanine can correspond, for example, to a level of about 400 pmol per 8×10^8 red blood cells. Where the

elevated 6-mercaptopurine metabolite is 6-thioguanine, the toxicity associated with 6-mercaptopurine treatment can be, for example, hematologic toxicity. The 6-mercaptopurine metabolite also can be a metabolite such as 6-methyl-mercaptopurine and the predetermined toxic level of 6-methyl-mercaptopurine can correspond, for example, to a level of about 7000 pmol per 8×10^8 red blood cells. Where the elevated 6-mercaptopurine metabolite is 6-methyl-mercaptopurine, the toxicity associated with 6-mercaptopurine treatment can be, for example, hepatic toxicity.

Further provided by the invention is a method of optimizing therapeutic efficacy and reducing toxicity associated with 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder. The method includes the steps of administering a 6-mercaptopurine drug to a subject having an immune-mediated gastrointestinal disorder; determining a level of 6-thioguanine in the subject having the immune-mediated gastrointestinal disorder; and determining a level of 6-methyl-mercaptopurine in the subject having the immune-mediated gastrointestinal disorder, where a level of 6-thioguanine less than a predetermined minimal therapeutic level indicates a need to increase the amount of 6-mercaptopurine drug subsequently administered to the subject, thereby increasing therapeutic efficacy; where a level of 6-thioguanine greater than a predetermined toxic level of 6-thioguanine indicates a need to decrease the amount of 6-mercaptopurine drug subsequently administered to the subject, thereby reducing toxicity associated with 6-mercaptopurine treatment of the immune-mediated gastrointestinal disorder; and where a level of 6-methyl-mercaptopurine greater than a predetermined toxic level of 6-methyl-mercaptopurine indicates a need

to decrease the amount of 6-mercaptopurine drug subsequently administered to the subject, thereby reducing toxicity associated with 6-mercaptopurine treatment of the immune-mediated gastrointestinal disorder.

In such a method of optimizing therapeutic efficacy and reducing toxicity associated with 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder, the predetermined minimal therapeutic level of 6-thioguanine can be, for example, a level corresponding to about 230 pmol per 8×10^8 red blood cells; the predetermined toxic level of 6-thioguanine can be, for example, a level corresponding to about 400 pmol per 8×10^8 red blood cells; and the predetermined toxic level of 6-methyl-mercaptopurine can be, for example, a level corresponding to about 7000 pmol per 8×10^8 red blood cells. The level of 6-thioguanine and 6-methyl-mercaptopurine each can be conveniently determined, for example, in red blood cells using high pressure liquid chromatography. The invention further provides methods to optimize the therapeutic efficacy of 6-mercaptopurine drug treatment of a non-IBD autoimmune disease.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows mercaptopurine metabolism and the structures of 6-mercaptopurine (6-MP) metabolites. The initial metabolism of 6-MP is catalyzed by thiopurine methyltransferase (TPMT), xanthine oxidase (XO), and hypoxanthine phosphoribosyltransferase (HPRT). Further metabolism of the thionucleotide is catalyzed by inosine monophosphate dehydrogenase (IMPD) and guanosine

monophosphate synthetase (GMPS). The breakdown of azathioprine to 6-mercaptopurine is nonenzymatic.

Figure 2 shows 6-mercaptopurine (6-MP) metabolism and the 6-MP metabolites that are measured as 6-MP, 6-thioguanine (6-TG) and 6-methyl-mercaptopurine (6-MMP) (indicated as "MP," "TG" and "MMP" inside the base). 6-TG mono-phosphate is converted to the di- and tri-phosphate by monophosphate kinase (MPK) and diphosphate kinase (DPK), respectively. The ribonucleoside diphosphate is converted to deoxyribonucleoside diphosphate by ribonucleotide reductase (RR).

Figure 3 shows ranges of 6-thioguanine in IBD patients treated with a 6-MP drug.

15 DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method of optimizing therapeutic efficacy of 6-mercaptopurine (6-MP) drug treatment of an immune-mediated gastrointestinal disorder. The method includes the steps of administering a 6-MP drug to a subject having an immune-mediated gastrointestinal disorder; and determining a level of 6-thioguanine (6-TG) in the subject having the immune-mediated gastrointestinal disorder, where a level of 6-TG less than a level corresponding to about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of 6-MP drug subsequently administered to the subject and where a level of 6-TG greater than a level corresponding to about 400 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of 6-MP drug subsequently administered to the subject. The methods are directed to

treating immune-mediated gastrointestinal disorders, including inflammatory bowel diseases (IBD) such as Crohn's disease and ulcerative colitis, lymphocytic colitis, microscopic colitis, collagenous colitis, 5 autoimmune enteropathy, allergic gastrointestinal disease and eosinophilic gastrointestinal disease. In a method of optimizing therapeutic efficacy of 6-MP treatment of IBD, the subject having IBD can be, for example, a pediatric subject. The level of 6-TG can be determined, 10 for example, in red blood cells using high pressure liquid chromatography (HPLC).

The invention provides methods of optimizing therapeutic efficacy of 6-MP drug treatment of an immune-mediated gastrointestinal disorder. The methods 15 of the invention are particularly useful for treating an immune-mediated gastrointestinal disorder such as IBD, including Crohn's disease and ulcerative colitis and subtypes thereof. The methods of the invention allow the clinician to provide an individually optimized dosage of 20 a 6-MP drug so as to achieve a target level of a 6-MP metabolite in a particular patient having an immune-mediated gastrointestinal disorder, thereby optimizing the effectiveness of 6-MP drug therapy in the patient. The methods of the invention for optimizing 25 therapeutic efficacy of 6-MP drug treatment involve determining the level of 6-TG in a patient having an immune-mediated gastrointestinal disorder. As disclosed herein, the level of 6-TG measured in a patient treated with a 6-MP drug was an indicator of the effectiveness of 30 drug treatment. A level of at least 230 pmol 6-TG/ 8×10^8 red blood cells (RBC) was found in responders to drug therapy (see Examples I and II). These results indicate that determining the level of 6-TG can be used to assess whether a patient has a level of 6-TG that is sufficient

to alleviate symptoms of an immune-mediated gastrointestinal disorder such as IBD, thus optimizing therapeutic efficacy.

As used herein, the term "6-mercaptopurine drug" or "6-MP drug" refers to any drug that can be metabolized to an active 6-mercaptopurine metabolite that has therapeutic efficacy such as 6-TG. Exemplary 6-mercaptopurine drugs as defined herein include 6-mercaptopurine (6-MP) and azathioprine (AZA). As illustrated in Figure 1, both of 6-MP and AZA can be metabolized to 6-mercaptopurine metabolites such as the exemplary 6-mercaptopurine metabolites shown in Figure 1, including 6-thioguanine (6-TG), 6-methyl-mercaptopurine (6-MMP) and 6-thiouric acid. (Lennard, Eur. J. Clin. Pharmacol. 43:329-339 (1992)).

Other 6-MP drugs include, for example, 6-methylmercaptopurine riboside and 6-TG (Loo et al., Clin. Pharmacol. Ther. 9:180-194 (1968); O'Dwyer et al., J. Natl. Cancer Inst. 83:1235-1240 (1991); Erb et al., Cancer Chemother. Pharmacol. 42:266-272 (1998); Lancaster et al., Br. J. Haematol. 102:439-443 (1998); Ingle et al., Am. J. Clin. Oncol. 20:69-72 (1997); Evans and Relling, Leuk. Res. 18:811-814 (1994)). 6-TG is a particularly useful 6-MP drug in patients having high TPMT activity. Patients exhibiting high TPMT activity are expected to more easily convert 6-MP drugs such as 6-MP and AZA to 6-MMP (see Figures 1 and 2). As disclosed herein, high levels of 6-MMP are associated with hepatotoxicity (see Examples I and II). Therefore, patients with high TPMT activity can be more susceptible to toxic effects of 6-MP drug therapy. By administering 6-TG, which is an active 6-MP metabolite associated with therapeutic efficacy (see Examples I and II), the

toxicity that can be associated with conversion of 6-MP to 6-MMP is bypassed.

It is understood that the 6-MP metabolites can be the metabolites shown in Figure 1 or analogues thereof. As used herein, the term "6-thioguanine" or "6-TG" refers to 6-thioguanine or analogues thereof, including molecules having the same base structure, for example, 6-thioguanine ribonucleoside, 6-thioguanine ribonucleotide mono-, di- and tri-phosphate, 6-thioguanine deoxyribonucleoside and 6-thioguanine deoxyribonucleotide mono-, di, and triphosphate. The term "6-TG" also includes derivatives of 6-thioguanine, including chemical modifications of 6-TG, so long as the structure of the 6-TG base is preserved.

As used herein, the term "6-methyl-mercaptopurine" or "6-MMP" refers to 6-methyl-mercaptopurine or analogues thereof, including analogues having the same base structure, for example, 6-methyl-mercaptopurine ribonucleoside, 6-methyl-mercaptopurine ribonucleotide mono-, di-, and tri-phosphate, 6-methyl-mercaptopurine deoxyribonucleoside, and 6-methyl-mercaptopurine deoxyribonucleotide mono-, di- and tri-phosphate. The term "6-MMP" also includes derivatives of 6-methyl-mercaptopurine, including chemical modifications of 6-MMP, so long as the structure of the 6-MMP base is preserved.

The methods of the invention relate to treatment of an immune-mediated gastrointestinal disorder. As used herein, the term "immune-mediated gastrointestinal disorder" or "immune-mediated GI disorder" refers to a non-infectious disease of the

gastrointestinal tract or bowel that is mediated by the immune system or cells of the immune system. Immune-mediated gastrointestinal disorders include, for example, inflammatory bowel diseases (IBD) such as Crohn's disease
5 and ulcerative colitis, lymphocytic colitis, microscopic colitis, collagenous colitis, autoimmune enteropathy, allergic gastrointestinal disease and eosinophilic gastrointestinal disease.

The methods of the invention are particularly
10 useful for treating IBD, or subtypes thereof, which has been classified into the broad categories of Crohn's disease and ulcerative colitis. As used herein, "a subject having inflammatory bowel disease" is synonymous with the term "a subject diagnosed with having an
15 inflammatory bowel disease," and means a patient having Crohn's disease or ulcerative colitis. Crohn's disease (regional enteritis) is a disease of chronic inflammation that can involve any part of the gastrointestinal tract. Commonly, the distal portion of the small intestine
20 (ileum) and cecum are affected. In other cases, the disease is confined to the small intestine, colon or anorectal region. Crohn's disease occasionally involves the duodenum and stomach, and more rarely the esophagus and oral cavity.

25 The variable clinical manifestations of Crohn's disease are, in part, a result of the varying anatomic localization of the disease. The most frequent symptoms of CD are abdominal pain, diarrhea and recurrent fever. CD is commonly associated with intestinal obstruction or
30 fistula, which is an abnormal passage between diseased loops of bowel, for example. Crohn's disease also includes complications such as inflammation of the eye, joints and skin; liver disease; kidney stones or

amyloidosis. In addition, CD is associated with an increased risk of intestinal cancer.

Several features are characteristic of the pathology of Crohn's disease. The inflammation associated with CD, known as transmural inflammation, involves all layers of the bowel wall. Thickening and edema, for example, typically also appear throughout the bowel wall, with fibrosis also present in long-standing disease. The inflammation characteristic of CD also is discontinuous in that segments of inflamed tissue, known as "skip lesions," are separated by apparently normal intestine. Furthermore, linear ulcerations, edema, and inflammation of the intervening tissue lead to a "cobblestone" appearance of the intestinal mucosa, which is distinctive of CD.

A hallmark of Crohn's disease is the presence of discrete aggregations of inflammatory cells, known as granulomas, which are generally found in the submucosa. Some Crohn's disease cases display the typical discrete granulomas, while others show nonspecific transmural inflammation. As a result, the presence of discrete granulomas is indicative of CD, although the absence of granulomas also is consistent with the disease. Thus, transmural or discontinuous inflammation, rather than the presence of granulomas, is a preferred diagnostic indicator of Crohn's disease (Rubin and Farber, Pathology (Second Edition) Philadelphia: J.B. Lippincott Company (1994)).

Ulcerative colitis (UC) is a disease of the large intestine characterized by chronic diarrhea with cramping abdominal pain, rectal bleeding, and loose discharges of blood, pus and mucus. The manifestations

of ulcerative colitis vary widely. A pattern of exacerbations and remissions typifies the clinical course of most UC patients (70%), although continuous symptoms without remission are present in some patients with UC.

- 5 Local and systemic complications of UC include arthritis, eye inflammation such as uveitis, skin ulcers and liver disease. In addition, ulcerative colitis and especially long-standing, extensive disease is associated with an increased risk of colon carcinoma.

- 10 Several pathologic features characterize UC in distinction to other inflammatory bowel diseases. Ulcerative colitis is a diffuse disease that usually extends from the most distal part of the rectum for a variable distance proximally. The term left-sided
- 15 colitis describes an inflammation that involves the distal portion of the colon, extending as far as the splenic flexure. Sparing of the rectum or involvement of the right side (proximal portion) of the colon alone is unusual in ulcerative colitis. The inflammatory process
- 20 of ulcerative colitis is limited to the colon and does not involve, for example, the small intestine, stomach or esophagus. In addition, ulcerative colitis is distinguished by a superficial inflammation of the mucosa that generally spares the deeper layers of the bowel
- 25 wall. Crypt abscesses, in which degenerated intestinal crypts are filled with neutrophils, also are typical of ulcerative colitis (Rubin and Farber, *supra*, 1994).

- In comparison with Crohn's disease, which is a patchy disease with frequent sparing of the rectum,
- 30 ulcerative colitis is characterized by a continuous inflammation of the colon that usually is more severe distally than proximally. The inflammation in ulcerative colitis is superficial in that it is usually limited to

the mucosal layer and is characterized by an acute inflammatory infiltrate with neutrophils and crypt abscesses. In contrast, Crohn's disease affects the entire thickness of the bowel wall with granulomas often, although not always, present. Disease that terminates at the ileocecal valve, or in the colon distal to it, is indicative of ulcerative colitis, while involvement of the terminal ileum, a cobblestone-like appearance, discrete ulcers or fistulas suggest Crohn's disease.

- 10 In addition to IBD, immune-mediated GI disorders also include other gastrointestinal diseases such as lymphocytic colitis; microscopic colitis; collagenous colitis; autoimmune enteropathy, including autoimmune enteritis and autoimmune enterocolitis; 15 allergic gastrointestinal disease; and eosinophilic gastrointestinal disease, including eosinophilic gastroenteritis and eosinophilic enteropathy.

- Over the past two decades, the histological evaluation of colorectal biopsies obtained by colonoscopy 20 has expanded the spectrum of chronic IBD. A new group of immune-mediated bowel disorders has emerged, characterized by chronic watery diarrhea, minimal or absent endoscopic findings, and inflammatory changes in mucosal biopsies. Lymphocytic colitis, also commonly 25 referred to as microscopic colitis, is a clinicopathological syndrome characterized primarily by lymphocytic infiltration of the epithelium. Collagenous colitis is defined by the presence of a collagenous band below the surface epithelium, accompanied by an increase 30 in inflammatory cell infiltrate (Lazenby et al. Hum. Pathol. 20:18-28 (1989)). These disorders are often associated with other autoimmune diseases such as rheumatoid arthritis, pernicious anemia, thyroiditis,

uveitis and type I diabetes mellitus. Clinicians have used immunosuppressive drugs, including 6-MP, to treat these disorders (Deslandres et al. J. Pediatr. Gastroenterol. Nutr. 25:341-346 (1997)).

- 5 Autoimmune enteropathy, including autoimmune enteritis and autoimmune enterocolitis, is a syndrome of severe secretory diarrhea and marked enterocolitis, in association with diagnostic circulating antibodies to enterocytes (Seidman et al., J. Pediatr. 117:929-932 (1990)). This syndrome, most often seen in infancy, can be seen in association with other autoimmune diseases. Complete villous atrophy is associated with a severe inflammatory reaction on small bowel biopsies. Although some cases remit after an extended period of time, most
- 10 patients die without immunosuppressive therapy, which can include 6-MP drug therapy.

- Eosinophilic gastrointestinal disease, including eosinophilic gastroenteritis and eosinophilic enteropathy, is characterized by a dense infiltration of eosinophils in one or more areas of the gastrointestinal tract, variable intestinal symptoms, and usually a peripheral eosinophilia (80% of cases). Food allergic, including allergic gastrointestinal disease, and eosinophilic disorders of the gastrointestinal tract are
- 20 commonly treated by dietary elimination of the offending nutrients. However, both food induced and eosinophilic enteropathies may, in certain circumstances, require corticosteroid and immunosuppressive therapy, including 6-MP (Russo et al., Pediatric Dev. Path. 2:65-71 (1999)).
- 30

The methods of the invention relate to optimizing therapeutic efficacy of 6-MP drug treatment of an immune-mediated GI disorder, including IBD such as Crohn's disease and ulcerative colitis and subtypes thereof. The methods of the invention are particularly useful for treating patients dependent on steroid therapy for maintenance of remission of disease in Crohn's disease and ulcerative colitis patients. As used herein, the phrase "optimizing therapeutic efficacy of 6-MP drug treatment" refers to adjusting the therapeutic dosage of a 6-MP drug such as 6-MP or azathioprine so that the concentration of a 6-MP metabolite that is correlated with effective treatment is maintained. As set forth above, the methods of the invention allow the clinician to provide an individually optimized dosage of a 6-MP drug so as to achieve a target level of a 6-MP metabolite in a particular patient, thereby optimizing the effectiveness of 6-MP drug therapy in the patient. Therapeutic efficacy generally is indicated by alleviation of one or more signs or symptoms associated with the disease. In the case of immune-mediated GI disorders, in particular IBD, therapeutic efficacy is indicated by alleviation of one or more signs or symptoms associated with the disease, including, for example, joint pain, arthritis, arthralgia, anorexia, growth failure, fistula closure, abdominal pain, diarrhea, recurrent fever, anemia, weight loss, rectal bleeding, inflammation of the intestine, and loose discharges of blood, pus and mucus. Methods for determining therapeutic efficacy, in particular for treating IBD, are disclosed herein in Examples I and II.

Therapeutic efficacy can be readily determined by one skilled in the art as the alleviation of one or more signs or symptoms of the disease being treated. In

the case of IBD, patients can be analyzed using a Crohn's disease activity index (Best et al., Gastroenterology 70:439-444 (1976)). IBD patients can also be analyzed using a Harvey-Bradshaw index (HBI) (Harvey and Bradshaw, 5 Lancet 1:514 (1980)). The Harvey-Bradshaw index provides an analytical method for measuring signs or symptoms of Crohn's disease, including the signs or symptoms of general well-being, abdominal pain, number of liquid stools per day, abdominal mass, and complications such as 10 arthralgia, uveitis, erythema nodosum, aphthous ulcers, pyoderma gangrenosum, anal fissure, new fistula and abscess. The Harvey-Bradshaw index is particularly useful when evaluating pediatric patients.

Previous studies suggested that measurement of 15 6-MP metabolite levels can be used to predict clinical efficacy and tolerance to azathioprine or 6-MP (Cuffari et al., Gut 39:401-406 (1996a)). However, it was unknown what concentrations of 6-MP metabolites correlated with optimized therapeutic efficacy or with toxicity (Cuffari 20 et al., *supra*, 1996a). As disclosed herein, levels of 6-MP metabolites such as 6-TG and 6-MMP were determined and correlated with therapeutic efficacy and toxicity associated with 6-MP drug therapy (see Examples I and II).

25 The invention is directed to methods of optimizing therapeutic efficacy of 6-MP drug treatment of an immune-mediated GI disorder by monitoring predetermined levels associated with therapeutic efficacy or toxicity and adjusting the 6-MP drug dosage so as to 30 maintain an optimized dose that is efficacious and has reduced toxicity. The methods involve administering a 6-MP drug such as 6-MP or azathioprine to a subject having an immune-mediated GI disorder and determining a

level of a 6-MP metabolite in the subject having the immune-mediated GI disorder. The methods of the invention are advantageous in that the dosage of a 6-MP drug can be adjusted to maximize the efficacy of treating
 5 an immune-mediated GI disorder such as IBD while minimizing toxicity associated with 6-MP drug treatment.

As used herein, the term "6-mercaptopurine metabolite" refers to a product derived from 6-mercaptopurine in a biological system. Exemplary
 10 6-mercaptopurine metabolites are shown in Figure 1 and include 6-thioguanine (6-TG), 6-methyl-mercaptopurine (6-MMP) and 6-thiouric acid and analogues thereof. For example, 6-MP metabolites include 6-TG bases such as 6-TG, 6-thioguanosine mono-, di- and tri-phosphate; 6-MMP
 15 bases such as 6-methyl-mercaptopurine and 6-methyl-thioinosine monophosphate; 6-thioxanthosine (6-TX) bases such as 6-thioxanthosine mono-phosphate; 6-thioruric acid (6-TUA); and 6-MP bases such as 6-mercaptopurine and 6-thioinosine monophosphate (see
 20 Figure 2). The immunosuppressive properties of 6-MP are believed to be mediated via the intracellular transformation of 6-MP to its active metabolites such as 6-TG and 6-MMP nucleotides. Furthermore, 6-MP metabolites such as 6-TG and 6-MMP were found to
 25 correlate with therapeutic efficacy and toxicity associated with 6-MP drug treatment of IBD patients (see Examples I and II).

The level of a 6-MP metabolite can be determined by methods well known in the art including,
 30 for example, those described in Lilleyman and Lennard, Lancet 343:1188-1190 (1994); Lennard and Singleton, J. Chromatography Biomed. Applicat. 583:83-90 (1992); Lennard and Singleton, J. Chromatography 661:25-33

(1994); and Cuffari et al., Can. J. Physiol. Pharmacol. 74:580-585 (1996b)). 6-MP metabolites such as 6-TG and 6-MMP can be measured, for example, by collecting red blood cells and extracting thiobases, for example, 6-MP, 5 6-TG, 6-TX and 6-MMP, which are released by acid hydrolysis. 6-MMP is converted to a form extractable by phenyl mercury salts (Dervieux and Boulieu, Clin. Chem. 44:2511-2525 (1998); Duchesne et al., Proc. Amer. Soc. Clin. Oncol. 13:137 (1994a); Duchesne et al., Can. J. 10 Physiol. Pharmacol. 72:197 (1994b)). Such an analysis measures the thiobase and its analogues, including ribonucleoside, ribonucleotide, deoxyribonucleoside, deoxyribonucleotide thiobases as well as mono-, di- and tri-phosphate analogues, which have been converted to 15 thiobases.

Acid hydrolyzed extracts can be analyzed by resolving 6-MP metabolites and measuring their levels. For example, HPLC such as reverse phase HPLC is a useful method for resolving and measuring the levels of 6-MP 20 metabolites, including 6-MP, 6-TG and 6-MMP (Lennard and Singleton, *supra*, 1992). Ultraviolet light (UV) detection can be used to measure the 6-MP metabolites. A particularly useful method of measuring 6-MP metabolites is isocratic reverse phase HPLC with UV detection 25 (Cuffari et al., *supra*, 1996b).

Other methods for measuring 6-MP metabolites can also be used. For example, ion-pairing HPLC with dual UV-wavelength detection can be used to measure 6-MP metabolites (Zimm and Strong, Anal. Biochem. 160:1-6 30 (1987)). Additional methods for measuring 6-MP metabolites include, for example, capillary electrophoresis with laser-induced fluorescence detection (Rabel et al., Anal. Biochem. 224:315-322 (1995)); anion

- exchange chromatography and fluorescent detection (Tidd and Dedhar J. Chromatography 145:237-246 (1978)); lanthanum precipitation, acid hydrolysis, back extraction and fluorometric assay (Fletcher and Maddocks, Brit. J. Clin. Pharmacol. 10:287-292 (1980)); thin layer chromatography (Bennet and Allen, Cancer Res. 31:152-158 (1971)); precolumn derivatization with the thiol-reactive fluorophore monobromobimane, treatment with alkaline phosphatase, HPLC resolution and quantification by fluorometry (Warren and Slordal, Anal. Biochem. 215:278-283 (1993)); and enzymatic hydrolysis followed by HPLC separation and UV detection (Giverhaug et al., Ther. Drug Monit. 19:663-668 (1997)). 6-MP metabolites such as 6-TG can also be measured in DNA by degrading DNA to deoxyribonucleosides, derivatizing deoxy-6-TG with a fluorophore and resolving on reverse phase HPLC (Warren et al., Cancer Res. 55:1670-1674 (1995)).

- As used herein, the level of a 6-MP metabolite can include the 6-MP metabolite itself, or the level of the 6-MP metabolite and analogues thereof. For example, as described above, acid hydrolysis can be used to release thionucleotides from a sample, resulting in conversion of mono-, di- and tri-phosphates to thiobases. In such an analysis, the level of several analogues of a thionucleotide are measured (see Figure 2). For example, measuring 6-TG can include 6-thioguanosine 5'-mono-, di-, and tri-phosphate, and 6-thiodeoxyguanosine 5'-di-, and tri-phosphate. Measuring 6-MP can include 6-mercaptopurine and 6-thioinosine 5'-monophosphate.
- Measuring 6-MMP can include 6-methylmercaptopurine and 6-methylthioinosine 5'-monophosphate, and can also include 6-methylthioinosine di- and tri-phosphate, as well as 6-methyl thioguanosine.

A particularly useful determination of the level of a 6-MP metabolite is the median level of the 6-MP metabolite since the distribution of 6-MP metabolite values is trimodal. Unless otherwise designated, the levels referred to herein are median levels. A 6-MP metabolite level can also be a mean level, if so desired. Unless otherwise designated, the levels of 6-MP metabolites referred to herein are values per 8×10^8 RBC, whether reported as mean or median values.

6-MP metabolite levels can be conveniently assayed using red blood cells because such cells are readily available from the patient, lack a nucleus and are easy to manipulate. However, it should be understood that any measurement that allows determination of 6-MP metabolite levels can be used. For example, leukocytes can be used to measure 6-MP metabolite levels, which can be correlated with 6-MP metabolite levels in erythrocytes (Cuffari et al., *supra*, 1996b). Regardless of the method employed to measure 6-MP metabolites, one skilled in the art can readily measure 6-MP metabolite levels in a sample, for example, in leukocytes or DNA obtained from a patient, and correlate the level of 6-MP metabolites to the values disclosed herein, which were determined using RBC.

For convenience, levels of 6-MP metabolites disclosed herein are given in terms of an assay with RBC and, therefore, are given as an amount of a 6-MP metabolite in a given number of RBCs. 6-MP metabolites assayed in RBCs can also be determined relative to the amount of hemoglobin. However, it should be understood that one skilled in the art can measure 6-MP metabolite levels in samples other than RBCs and readily correlate such levels with 6-MP metabolite levels in RBCs. For

example, one skilled in the art can readily determine levels of a 6-MP metabolite in cells such as leukocytes, or cells from the oral mucosa, and in RBCs by measuring the level of 6-MP metabolites in both types of cells and
5 determining the correspondence between levels of 6-MP metabolites in RBCs and the levels in the other cells. Once a correspondence between 6-MP metabolites in RBCs and in a sample has been determined, one skilled in the art can use that correspondence to measure levels in the
10 other sample and correlate those levels with the levels in RBCs disclosed herein.

As disclosed herein, the level of 6-TG in an IBD patient treated with a 6-MP drug was found to correlate with therapeutic efficacy (see Examples I and
15 II). In particular, a median level of 230 pmol 6-TG/ 8×10^8 RBC or more was found in IBD patients who responded to 6-MP drug therapy. Thus, as disclosed herein, a level of at least about 230 pmol 6-TG per 8×10^8 RBC can be a minimal therapeutic level of 6-TG for efficaciously
20 treating a patient. Accordingly, a level of 6-TG below about 230 pmol/ 8×10^8 RBC indicates a need to increase the amount of a 6-MP drug administered to the patient. A minimal therapeutic level of 6-TG for efficaciously treating a patient also can be, for example, at least
25 about 240 pmol per 8×10^8 RBC; at least about 250 pmol per 8×10^8 RBC; at least about 260 pmol per 8×10^8 RBC; at least about 280 mol per 8×10^8 RBC or at least about 300 pmol per 8×10^8 RBC. It is understood that the minimal therapeutic levels of 6-TG disclosed herein are useful for treating
30 immune-mediated gastrointestinal disorders, including IBD, as well as non-IBD autoimmune diseases.

The methods of the invention directed to determining whether a patient has a minimal therapeutic level of a 6-MP metabolite are useful for indicating to the clinician a need to monitor a patient for therapeutic efficacy and to adjust the 6-MP drug dose, as desired. For example, in a patient having less than a minimal therapeutic level of a 6-MP metabolite such as 6-TG and who also presents as unresponsive to 6-MP drug therapy or having poor responsiveness to 6-MP drug therapy as measured by minimal or no effect on a sign or symptom of the disease being treated, one skilled in the art can determine that the dosage of a 6-MP drug should be increased. However, if it is determined that a patient has less than a predetermined minimal therapeutic level of a 6-MP metabolite but is responsive to 6-MP therapy, the current dose of 6-MP drug can be maintained. Based on measuring 6-MP metabolite levels and assessing the responsiveness of the patient to 6-MP therapy, one skilled in the art can determine whether a 6-MP drug dose should be maintained, increased, or decreased.

Although 6-MP drugs such as 6-MP and azathioprine can be used for effective treatment of an immune-mediated GI disorder, including IBDs such as Crohn's disease or ulcerative colitis, administration of such drugs can be associated with toxic side effects. Toxicities associated with 6-MP drug administration include pancreatitis, bone marrow depression, allergic reactions and drug hepatitis as well as neoplasms and infectious complications (Present et al., Annals Int. Med. 111:641-649 (1989); Cuffari et al., *supra*, 1996a). As disclosed herein, various toxicities associated with 6-MP drug treatment, including hepatic toxicity, pancreatic toxicity and hematologic toxicity, correlate

with the level of 6-MP metabolites in a subject administered a 6-MP drug (see Examples I and II).

Thus, the present invention also provides a method of reducing toxicity associated with 6-MP drug treatment of an immune-mediated GI disorder. The method of the invention for reducing toxicity associated with 6-MP drug treatment includes the steps of administering a 6-MP drug to a subject having an immune-mediated GI disorder; and determining a level of a 6-MP metabolite in the subject having the immune-mediated GI disorder, where a level of the 6-MP metabolite greater than a predetermined toxic level of the 6-MP metabolite indicates a need to decrease the amount of 6-MP drug subsequently administered to the subject, thereby reducing toxicity associated with 6-MP drug treatment of the immune-mediated GI disorder. In a method of the invention, the 6-MP metabolite can be, for example, 6-TG and the predetermined toxic level of 6-TG can correspond, for example, to a level of about 400 pmol per 8×10^8 red blood cells. Where the elevated 6-MP metabolite is 6-TG, the toxicity associated with 6-MP treatment can be, for example, hematologic toxicity, including leukopenia or bone marrow suppression. The 6-MP metabolite also can be a metabolite such as 6-MMP, and the predetermined toxic level of 6-MMP can correspond, for example, to a level of about 7000 pmol per 8×10^8 red blood cells. Where the elevated 6-MP metabolite is 6-MMP, the toxicity associated with 6-MP drug treatment can be, for example, hepatic toxicity.

As disclosed herein, the level of a 6-MP metabolite can be determined in a subject treated with a 6-MP drug and compared to a predetermined toxic level of a 6-MP metabolite such as 6-TG or 6-MMP to adjust future

6-MP drug administration, thereby reducing toxicity in the subject. For example, as disclosed herein, levels of 6-TG above about 400 pmol/ 8×10^8 RBC indicated that a patient was likely to experience toxicity, in particular hematologic toxicity such as leukopenia (see Examples I and II). Accordingly, a level of 6-TG above about 400 pmol/ 8×10^8 RBC can be a predetermined toxic level of 6-TG, which indicates that the amount of 6-MP drug subsequently administered should be decreased.

It is understood that, when a patient is determined to have a level of a 6-MP metabolite such as 6-TG or 6-MMP higher than a predetermined toxic level, one skilled in the art can make a determination as to whether a 6-MP drug dose should be decreased. For example, if the level of a 6-MP metabolite such as 6-TG or 6-MMP is higher than a predetermined toxic level, one skilled in the art can monitor for toxic side effects by measuring one or more of the toxicities associated with 6-MP drug treatment, as disclosed herein. As disclosed herein, a level of 6-TG greater than about 400 pmol per 8×10^8 RBCs was associated with increased risk of leukopenia or bone marrow suppression. Therefore, one skilled in the art can measure white blood cells (WBC) in a patient having levels of 6-TG higher than a predetermined toxic level to determine if the patient is exhibiting signs of reduced WBC counts. If such a patient exhibits signs of leukopenia or bone marrow suppression, the 6-MP drug dose can be reduced. However, if it is determined that a patient has levels of a 6-MP metabolite higher than a predetermined toxic level but does not exhibit signs of leukopenia or other 6-MP drug toxicities, one skilled in the art can determine that the current 6-MP drug dose can be maintained. Based on measuring 6-MP metabolite levels and determining signs or

symptoms of toxicities associated with 6-MP drug treatment, one skilled in the art can determine whether a 6-MP drug dose should be maintained or decreased. As such, a level of a 6-MP metabolite higher than a predetermined toxic level can indicate a need to measure a toxicity associated with 6-MP drug treatment such as measuring WBCs or any of the other signs or symptoms of toxicities associated with 6-MP drug treatment to determine if the 6-MP drug dose should be adjusted.

Furthermore, it is understood that, when decreasing the 6-MP drug dose, one skilled in the art will know or can readily determine whether the 6-MP drug dose should be decreased to a lower dose or whether 6-MP drug administration should be stopped for some period of time, or terminated. For example, if the clinician determines that 6-MP drug therapy should be stopped for some period of time due to levels of a 6-MP metabolite exceeding a level predetermined to be toxic, the levels of 6-MP metabolites can be monitored after stopping 6-MP drug therapy until the level of the toxic 6-MP metabolite returns to a safe, non-toxic level. At that time, the clinician can resume 6-MP drug therapy, if desired.

The methods of the invention for reducing toxicity associated with 6-MP drug treatment of a disease involve comparing a level of a 6-MP metabolite to a predetermined toxic level of a 6-MP metabolite. In general, a "predetermined toxic level" of a 6-MP metabolite means a level of a 6-MP metabolite that has been correlated with one or more toxicities associated with 6-MP drug treatment. As disclosed herein, a predetermined toxic level of 6-TG can be about 400 pmol per 8×10^8 RBC. A predetermined toxic level of 6-TG also can be about 350 pmol per 8×10^8 RBC; 370 pmol per 8×10^8

RBC; 390 pmol per 8×10^8 RBC; 425 pmol per 8×10^8 RBC; or 450 pmol per 8×10^8 RBC. It is understood that the predetermined toxic levels of 6-TG disclosed herein are useful for treating immune-mediated GI disorders, including IBD, as well as non-IBD autoimmune diseases.

Another 6-MP metabolite useful for predicting the likelihood of toxicity is 6-methyl-mercaptopurine (6-MMP). As disclosed herein, a level of greater than about 7000 pmol 6-MMP/ 8×10^8 in patients administered a 6-MP drug was associated with toxicity, in particular hepatotoxicity (see Examples I and II). These results indicate that the level of 6-MMP can be used to predict toxicity in a patient treated with a 6-MP drug. As disclosed herein, a predetermined toxic level of 6-MMP can be about 7000 pmol per 8×10^8 RBC. A predetermined toxic level of 6-MMP also can be about 6000 pmol per 8×10^8 RBC; 6500 pmol per 8×10^8 RBC; 7500 pmol per 8×10^8 RBC; or 8000 pmol per 8×10^8 RBC. It is understood that the predetermined toxic levels of 6-MMP disclosed herein are useful for treating immune-mediated GI disorders, including IBD, as well as non-IBD autoimmune diseases. According to a method of the invention, if the level of 6-MMP is above a predetermined toxic level, the subsequent administration of a 6-MP drug can be decreased to minimize toxicity.

Further provided by the invention is a method of optimizing therapeutic efficacy and reducing toxicity associated with 6-MP drug treatment of an immune-mediated GI disorder such as IBD. The method includes the steps of administering a 6-MP drug to a subject having an immune-mediated GI disorder; determining a level of 6-TG in the subject having the immune-mediated GI disorder; and determining a level of 6-MMP in the subject having

the immune-mediated GI disorder, where a level of 6-TG less than a predetermined minimal therapeutic level indicates a need to increase the amount of 6-MP drug subsequently administered to the subject, thereby

5 increasing therapeutic efficacy; where a level of 6-TG greater than a predetermined toxic level of 6-TG indicates a need to decrease the amount of 6-MP drug subsequently administered to the subject, thereby

10 reducing toxicity associated with 6-MP treatment of the immune-mediated GI disorder; and where a level of 6-MMP greater than a predetermined toxic level of 6-MMP indicates a need to decrease the amount of 6-MP drug subsequently administered to the subject, thereby

15 reducing toxicity associated with 6-MP drug treatment of the immune-mediated GI disorder.

In such a method of optimizing therapeutic efficacy and reducing toxicity associated with 6-MP drug treatment of an immune-mediated GI disorder such as IBD, the predetermined minimal therapeutic level of 6-TG can

20 be, for example, a level corresponding to about 230 pmol per 8×10^8 red blood cells; the predetermined toxic level of 6-TG can be, for example, a level corresponding to about 400 pmol per 8×10^8 red blood cells; and the predetermined toxic level of 6-MMP can be, for example, a

25 level corresponding to about 7000 pmol per 8×10^8 red blood cells. In a method of the invention, the subject having an immune-mediated GI disorder such as IBD can be, for example, a pediatric subject. The level of 6-TG and 6-MMP each can be conveniently determined, for example,

30 in red blood cells using HPLC.

The methods of the invention are useful for optimizing the amount of a 6-MP drug to be administered to a patient with an immune-mediated GI disorder such as

IBD. By measuring the levels of 6-MP metabolites such as 6-MMP and 6-TG, one skilled in the art can determine the 6-MP drug dosage that will result in optimized therapeutic efficacy and reduced toxicity when administered to a patient.

As disclosed herein, gender and age differences were observed in pediatric patients treated with 6-MP drug therapy (see Example III). Very little change in 6-MP metabolite levels of 6-TG and 6-MMP was seen for girls who had gone through puberty (older than age 12). However, boys who had gone through puberty (older than age 14) had a marked decrease in the level of 6-MMP, suggesting that hormonal changes occurring during puberty can affect the metabolism of a 6-MP drug. Therefore, the methods of the invention can additionally be used to monitor 6-MP metabolite levels in adolescents, particularly those going through puberty, in order to optimize therapeutic efficacy or minimize toxic side effects associated with 6-MP therapy.

As disclosed herein, TPMT genotyping is useful for predicting the effectiveness of 6-MP therapy in an IBD patient (see Example IV). Heterozygote patients are expected to have lower TPMT activity and should therefore be monitored for high levels of 6-TG for possible toxic levels associated with leukopenia or bone marrow suppression. 6-MP drug doses can be reduced accordingly. Wild type homozygous patients are expected to have higher TPMT activity and should therefore be monitored to maintain an effective therapeutic level of 6-TG and to determine if patients develop toxic levels of 6-MMP. Homozygous patients deficient in TPMT activity can be treated with lower doses of a 6-MP drug provided that patients are closely monitored for toxicity such as

leukopenia. Therefore, TPMT genotyping can be used to predict patient responsiveness to and potential toxicities associated with 6-MP drug therapy.

Furthermore, TPMT genotyping can be combined with other
5 methods of the invention to both determine TPMT genotype and to monitor 6-MP metabolites. TPMT genotyping can be particularly valuable when determining a starting dose of 6-MP drug therapy but can also be useful when adjusting 6-MP drug doses after therapy has begun.

10 The invention additionally provides a method of optimizing therapeutic efficacy of 6-MP drug treatment of a non-IBD autoimmune disease. The method includes the steps of administering a 6-MP drug to a subject having a non-IBD autoimmune disease; and determining a level of
15 6-thioguanine (6-TG) in the subject having the non-IBD autoimmune disease, where a level of 6-TG less than a minimal therapeutic level indicates a need to increase the amount of 6-MP drug subsequently administered to the subject and where a level of 6-TG greater than a
20 predetermined toxic level indicates a need to decrease the amount of 6-MP drug subsequently administered to the subject. The level of 6-MMP can also be monitored in a patient having a non-IBD autoimmune disease to determine if the level is higher than a predetermined toxic level
25 of 6-MMP.

The methods of the invention can be used to optimize therapeutic efficacy of 6-MP drug treatment of a non-IBD autoimmune disease. Such a non-IBD autoimmune disease can be any non-IBD autoimmune disease treatable
30 by a 6-MP drug such as 6-MP or azathioprine and, in particular, can be a disease such as rheumatoid arthritis, systemic lupus erythematosus, autoimmune

hepatitis (chronic active hepatitis) or pemphigus vulgaris.

As used herein, the term "non-IBD autoimmune disease" means a disease resulting from an immune response against a self tissue or tissue component, including both self antibody responses and cell-mediated responses. The term non-IBD autoimmune disease encompasses organ-specific non-IBD autoimmune diseases, in which an autoimmune response is directed against a single tissue, including myasthenia gravis, vitiligo, Graves' disease, Hashimoto's disease, Addison's disease, autoimmune gastritis, and Type I diabetes mellitus. The term non-IBD autoimmune disease also encompasses non-organ specific autoimmune diseases, in which an autoimmune response is directed against a component present in several or many organs throughout the body. Non-organ specific autoimmune diseases include, for example, systemic lupus erythematosus, progressive systemic sclerosis and variants, polymyositis and dermatomyositis, and rheumatoid disease. Additional non-IBD autoimmune diseases include pernicious anemia, primary biliary cirrhosis, autoimmune thrombocytopenia, and Sjögren's syndrome. One skilled in the art understands that the methods of the invention can be applied to these or other non-IBD autoimmune diseases treatable by a 6-MP drug such as 6-MP or azathioprine, or other 6-MP drugs, as desired. Specifically excluded from the term "non-IBD autoimmune disease" are diseases resulting from a graft versus host response and inflammatory bowel diseases such as Crohn's disease or ulcerative colitis.

The methods of the invention are also useful for treating a non-immune-mediated GI disorder autoimmune disease. As used herein, the term "non-immune-mediated GI disorder autoimmune disease" is a non-IBD autoimmune disease and specifically excludes immune-mediated GI disorders.

The methods of the invention can be particularly useful for optimizing therapeutic efficacy of 6-MP drug treatment of rheumatoid arthritis.

10 Rheumatoid arthritis is a chronic systemic disease primarily of the joints, usually polyarticular, marked by inflammatory changes in the synovial membranes and articular structures and by muscle atrophy and rarefaction of the bones.

15 The methods of the invention also can be particularly valuable for optimizing therapeutic efficacy of 6-MP drug treatment of lupus erythematosus and, in particular, systemic lupus erythematosus (SLE). Systemic lupus erythematosus is a chronic, remitting, relapsing
20 inflammatory, and sometimes febrile multisystemic disorder of connective tissue. SLE can be acute or insidious at onset and is characterized principally by involvement of the skin, joints, kidneys and serosal membranes.

25 Autoimmune hepatitis, also called chronic active hepatitis, also can be treated with a 6-MP drug and the dose optimized using the methods of the invention. Autoimmune hepatitis is a chronic inflammation of the liver occurring as a sequel to
30 hepatitis B or non-A, non-B hepatitis and is characterized by infiltration of portal areas by plasma cells and macrophages, piecemeal necrosis, and fibrosis.

The methods of the invention also can be useful for treating pemphigus vulgaris, the most common and severe form of pemphigus, which is a chronic, relapsing and sometimes fatal skin disease characterized clinically
5 by the development of successive crops of vesicles and bullae and treated by azathioprine. This disorder is characterized histologically by acantholysis, and immunologically by serum autoantibodies against antigens in the intracellular zones of the epidermis.

10 The methods of the invention can also be used to optimize the therapeutic efficacy of 6-MP drug treatment of graft versus host disease, which can occur in transplant patients. Graft versus host disease occurs when a transplant patient has an immune reaction to the
15 non-self transplant organ or tissue. The methods of the invention for optimizing the therapeutic efficacy of 6-MP drug treatment is particularly useful for treating heart, kidney and liver transplant recipients. The methods of the invention can be used to optimize therapeutic
20 efficacy and/or minimize toxicity associated with 6-MP drug treatment of a transplant patient.

The following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

25 6-MERCAPTOPURINE METABOLITE LEVELS PREDICT CLINICAL EFFICACY AND DRUG TOXICITY IN PEDIATRIC IBD

This example describes measuring 6-MP metabolite levels and correlation with the response of IBD patients treated with a 6-MP drug.

The levels of the 6-MP metabolites 6-TG and 6-MMP were measured in IBD patients to whom 6-MP pro-drug was administered, and the relationship of 6-MP metabolites to clinical disease activity and drug toxicity was determined. Briefly, blood was sampled (n=89) prior to daily administration of 6-MP in 55 IBD patients (CD n=51, UC n=4) receiving 1-1.5 mg/kg/day over at least a 4 month period (≥ 4 mo.). When AZA was administered, a conversion factor of 2.07 was used to convert to the equivalent 6-MP dose. Twice as much AZA is administered relative to 6-MP to have an equivalent dose of 6-MP.

Erythrocyte 6-TG, 6-MMP and 6-MP thiobases were measured (pmol/ 8×10^8 RBC) using reverse phase HPLC.

Briefly, blood samples were collected in EDTA (ethylene diamine tetraacetic acid) as anticoagulant. Cells were centrifuged and washed three times with an equal volume of 0.9% saline. Washed packed cells were stored at -70°C until analysis was performed.

For acid hydrolysis, 500 μl deionized H_2O , 50 μl thawed red blood cells, 40 μl of the appropriate standard or control, 500 μl 3.0 N H_2SO_4 , and 300 μl 10 mM dithiothreitol (DTT) was added to an 8 ml glass screw cap tube. The capped tubes were placed in a heating block preheated to 100°C and hydrolyzed. For 6-MMP, hydrolysis was carried out for 5 hours. For 6-MP and 6-TG, hydrolysis was carried out for 1 hour. After the incubation, tubes were cooled in a room temperature water bath. To tubes hydrolyzed for 5 hours (6-MMP), 400 μl 3.4 N NaOH and 1.0 ml 2 M Tris buffer, pH 9.0 was added. To tubes hydrolyzed for 1 hour (6-MP/6-TG), 450 μl 3.4 N NaOH and 500 μl 2M Tris base was added. A volume of 4 ml 0.03% phenylmercuric chloride in methylene chloride was

added to each tube. The tubes were capped and lightly agitated on a bi-directional rotator (15 min for 6-MMP and 30 min for 6-MP/6-TG). The contents were transferred to a 15 ml polypropylene centrifuge tube and centrifuged
5 at 3500 rpm for 3 min at 10°C. The aqueous phase (top layer) was discarded, and 3.0 ml of the organic phase (bottom layer) was transferred to a clean 15 ml polypropylene centrifuge tube. The analytes in the organic phase were back extracted by adding 225 μ l 0.1 N
10 HCl and lightly mixing on an orbital rotator for 5 min. Following vortexing for 30 seconds, the tubes were centrifuged at 3500 rpm for 3 min at 10°C.

For 6-MMP analysis, 50 μ l analyte was analyzed on a C18 reverse phase column with the mobile phase
15 containing 1 mM DTT, 2.078% triethylamine and 4% methanol, adjusted to pH 3.2 with concentrated H_3PO_4 . For 6-MP and 6-TG analysis, 100 μ l analyte was analyzed on a C18 reverse phase column using 0.1 M H_3PO_4 and 1 mM DTT in H_2O as the mobile phase.

20 Hepatic, pancreatic and hematological tests were obtained every 3 months. Clinical remission was defined as a Harvey Bradshaw Index < 5 in those patients off corticosteroids or weaned to a level of prednisone \leq 0.4 mg/kg/od (administered every other day). Treatment
25 failures were defined as non-responders (HBI > 5 or steroid dependence) or cessation of 6-MP due to side effects.

As shown in Table 1, a 6-TG level of >225 pmol per 8×10^8 RBC was associated with remission. The median
30 values shown in the tables represent pmol of the indicated 6-MP metabolite per 8×10^8 RBC. Excessive 6-TG and 6-MMP levels were associated with leukopenia and

hepatotoxicity, respectively. Negligible metabolite levels detected non-compliance as a cause of treatment failure in 2/31 cases. These results indicate that 6-MP metabolite levels predict both clinical responsiveness and drug-related toxicity.

Table 1:

Group	n	median 6-TG	6-TG >225	median 6-MMP
Responders	58	295	45/58 (78%)	3094
Non-responders	31	184*	8/31 (26%)	2048
Hepatic toxicity	7	258	5/7	9211**
Pancreatic toxicity	6	211	2/6	2342
Hematologic toxicity	6	414+	5/5	7042
p Values		*<0.001	‡0<0.001	**<0.001
		+<0.03		

These results demonstrate that determining levels of 6-MP metabolites is useful for predicting efficacy and toxicity of 6-MP drug therapy administered to IBD patients.

EXAMPLE II

6-MERCAPTOPURINE METABOLITE LEVELS CORRELATE WITH OPTIMAL 6-MP THERAPY IN IBD PATIENTS

This example describes prospective examination of the correlation of 6-MP metabolite levels with therapeutic response to 6-MP drug therapy and 6-MP drug related toxicity in IBD patients treated with 6-MP.

To obtain additional statistical data on IBD patients treated with a 6-MP drug, additional patients and samples were analyzed and combined with the data obtained in Example I. Blood was sampled at least once

5 in 93 IBD patients followed at Sainte-Justine Hospital IBD Center, Montreal, Canada, who were administered 6-MP drug therapy for at least 4 months. The 93 patients were pediatric patients, with 80 diagnosed as having CD, 8 diagnosed as having UC, and 5 diagnosed as having

10 indeterminate colitis (CD or UC). All but 7 patients were given AZA. The dosages were converted to 6-MP equivalents using a factor of 2.07 as described in Example I. For some patients, two or more samples were obtained and analyzed. Response to 6-MP was defined by

15 clinical remission (HBI <5, closed fistula) without corticosteroids. Disease activity and physical exam were ascertained at each clinic visit at which 6-MP metabolite levels were determined (clinical evaluation point). Hematological, pancreatic and hepatic laboratory

20 parameters were evaluated simultaneously. Erythrocyte 6-TG and 6-MMP concentrations ($\text{pmol}/8 \times 10^8 \text{ RBC}$) were measured by HPLC (Cuffari et al., *supra*, 1996a).

The results of the analysis of 6-MP metabolites in IBD patients are shown in Table 2. The number of

25 samples corresponds to the number of different samples obtained from the 93 patients. 6-TG quartile analysis, in which values are determined at 25, 50 and 75% of the data set, revealed that the frequency of response significantly increased at levels $> 230 \text{ pmol}/8 \times 10^8 \text{ RBC}$

30 ($p < 0.01$). Among patients in relapse, only 28% of patients had 6-TG levels $> 230 \text{ pmol}/8 \times 10^8 \text{ RBC}$. In contrast, 65% of patients in remission had 6-TG levels $> 230 \text{ pmol}/8 \times 10^8 \text{ RBC}$ ($p < 0.01$). Therefore, erythrocyte 6-TG concentrations were significantly and independently

associated with therapeutic response to 6-MP drug therapy.

The induction and maintenance of remission was found to be optimal at 6-TG levels $>230 \text{ pmol}/8 \times 10^8 \text{ RBC}$.

- 5 78% of patients above a median 6-TG of $230 \text{ pmol}/8 \times 10^8 \text{ RBC}$ were responders (see Figure 3). These results indicate that a 6-TG value of $230 \text{ pmol}/8 \times 10^8 \text{ RBC}$ can be used to predict efficacy of drug treatment with a 6-MP drug such as 6-MP or azathioprine.

10 Table 2:

Clinical evaluation point	(n)	Median 6-TG	Median 6-MMP	Median dose (mg/kg/day)
Remission	106	309*	2600	1.3
15 Relapse	72	197	1602	1.25
*p value		<0.0001	0.3	0.4

- Toxicity in the IBD patients treated with a 6-MP drug was also evaluated. Toxicity was measured essentially as described previously (Cuffari et al.,
 20 *supra*, 1996a). Thirty six patients (39%) experienced an adverse event. Hepatotoxicity was observed in 17% of patients, measured as the serum level of alanine aminotransferase (ALT) or aspartate aminotransferase (AST) (ALT, AST; exceeding or greater than 2x upper limit
 25 of normal). Leukopenia was observed in 14% of patients (white blood cell (WBC) < 4000). Pancreatic toxicity was observed in 7% of patients (lipase/amylase $>2 \times \text{N}$). High 6-MMP levels correlated significantly with hepatotoxicity (5463 with hepatotoxicity versus 2177 without
 30 hepatotoxicity; $p=0.04$). Leukopenia was observed in only 8% (8/106) of samples from patients in remission, with

significantly higher 6-TG levels observed in these patients (mean value of 490 in patients with leukopenia versus mean value of 323 without leukopenia; $p < 0.04$; median values were 342 versus 307, respectively).

- 5 Therefore, leukopenia did not correlate with therapeutic efficacy. Furthermore, drug dose (per kg) did not correlate with therapeutic efficacy (see Table 2). However, those patients who do develop leukopenia have a higher 6-TG level than the rest of the responder group.
- 10 These results indicate that 6-TG levels should be monitored to avoid potential clinical bone marrow suppression in responder patients who have high levels of 6-TG.

- These results demonstrate a significant
- 15 correlation between erythrocyte 6-TG levels and the therapeutic response to 6-MP drug treatment in IBD patients. The induction and maintenance of remission was found to be optimal at 6-TG levels $>230 \text{ pmol}/8 \times 10^8 \text{ RBC}$. Therefore, monitoring 6-MP metabolite levels, in
 - 20 particular 6-TG, is useful for determining that a therapeutically effective concentration of 6-MP metabolites is maintained while treating IBD patients with a 6-MP drug. Monitoring 6-MP metabolite levels, in particular 6-TG and 6-MMP, is also useful for minimizing
 - 25 6-MP drug related toxicity.

EXAMPLE III

GENDER AND AGE DIFFERENCES IN METABOLISM OF A 6-MP DRUG

- This example describes gender and age differences observed in pediatric patients treated with
- 30 6-MP drug therapy.

Pediatric IBD patients undergoing 6-MP drug therapy were assessed for levels of 6-MP metabolites. These patients were wild type for TPMT. Patients were assessed based on gender and age as it relates to

5 puberty. Puberty is established at 12 years of age in girls and 14 years of age in boys.

As shown in Table 3, the 6-MMP values are much lower in boys after puberty (greater than 14 years). Since the total amount of thiometabolites is lower, this

10 indicates that either lower doses of 6-MP are used or that there is a difference in the bioavailability of 6-MP after puberty in males.

Table 3: 6-MP Metabolite Levels in Pediatric IBD Patients

	Number Observed	6-TG level (pmol per 8×10^8 RBC)	6-MMP level (pmol per 8×10^8 RBC)	Ratio 6-MMP/6-TG
15 Girls (0-12y)	39	182.2	3447.0	18.52
Girls (>12y)	116	217.2	3304.5	12.55
Boys (0-14y)	51	235.0	3681.6	17.24
20 Boys (>14y)	104	222.5	1662.3*	6.78*
*p<0.001				

These results demonstrate that gender and age can affect the metabolism of 6-MP in pediatric IBD

25 patients undergoing 6-MP drug therapy.

EXAMPLE IV**THIOPURINE METHYLTRANSFERASE (TPMT) GENOTYPING AND
RESPONSIVENESS TO 6-MP DRUG THERAPY**

This example describes TPMT genotyping of IBD
5 patients treated with 6-MP drug therapy.

The genotype of TPMT was determined in IBD
patients that were responders and non responders.
Genotyping of TPMT was measured essentially as described
previously (Baccichet et al., Leuk. Res. 21:817-823
10 (1997); Zietkiewicz et al., Gene 205:161-171 (1997)).
The data shown in Table 4 indicate that patients
heterozygous for the TPMT mutation had significantly
higher 6-TG levels compared to those patients without the
mutation. All heterozygote patients were responders to
15 6-MP.

Table 4: TPMT Genotyping of IBD Patients

	Heterozygote	Normal
Responder	100%	55%
Non responder	0%	45%
mean 6-TG	589*	247

20 p value *less than 0.0001

TPMT genotyping revealed that 8 of 93 (9%) of
patients were heterozygotes. No homozygous TPMT
deficient patients were detected. All 8 heterozygotes
responded to 6-MP and had 6-TG levels >230 pmol/8x10⁸ RBC.

25 These results demonstrate that TPMT genotyping
is useful for predicting the effectiveness of 6-MP
therapy in an IBD patient. Heterozygote patients are
expected to have lower TPMT activity and should therefore

be monitored for high levels of 6-TG for possible toxic levels associated with leukopenia or bone marrow suppression. 6-MP drug doses can be reduced accordingly. Wild type homozygous patients are expected to have higher
5 TPMT activity and should therefore be monitored to maintain an effective therapeutic level of 6-TG and to determine if patients develop toxic levels of 6-MMP. Homozygous patients deficient in TPMT activity can be treated with lower doses of a 6-MP drug provided that
10 patients are closely monitored for toxicity such as leukopenia.

All journal articles and references provided herein, in parenthesis or otherwise, are incorporated herein by reference.

15 Although the invention has been described with reference to the examples provided above, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the claims.

We claim:

1. A method of optimizing therapeutic efficacy of 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder, comprising:

5 (a) administering a 6-mercaptopurine drug to a subject having said immune-mediated gastrointestinal disorder; and

(b) determining a level of 6-thioguanine in said subject having said immune-mediated gastrointestinal disorder,
10

wherein a level of 6-thioguanine less than a level corresponding to about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of 6-mercaptopurine drug subsequently administered to said
15 subject and

wherein a level of 6-thioguanine greater than a level corresponding to about 400 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of 6-mercaptopurine drug subsequently administered to said
20 subject.

2. The method of claim 1, wherein said immune-mediated gastrointestinal disorder is inflammatory bowel disease (IBD).

3. The method of claim 2, wherein said
25 subject having IBD is a pediatric subject.

4. The method of claim 1, wherein said immune-mediated gastrointestinal disorder is selected from the group consisting of lymphocytic colitis, microscopic colitis, collagenous colitis, autoimmune enteropathy, allergic gastrointestinal disease and eosinophilic gastrointestinal disease.

5. The method of claim 1, wherein said level of 6-thioguanine is determined in red blood cells.

6. The method of claim 5, wherein said level is determined using high pressure liquid chromatography.

7. A method of reducing toxicity associated with 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder, comprising:

(a) administering a 6-mercaptopurine drug to a subject having said immune-mediated gastrointestinal disorder; and

(b) determining a level of a 6-mercaptopurine metabolite in said subject having said immune-mediated gastrointestinal disorder,

20

wherein a level of said 6-mercaptopurine metabolite greater than a predetermined toxic level of said 6-mercaptopurine metabolite indicates a need to decrease the amount of 6-mercaptopurine drug subsequently administered to said subject, thereby reducing toxicity associated with 6-mercaptopurine drug treatment of said immune-mediated gastrointestinal disorder.

8. The method of claim 7, wherein said immune-mediated gastrointestinal disorder is IBD.

9. The method of claim 8, wherein said subject having IBD is a pediatric subject.

10. The method of claim 7, wherein said immune-mediated gastrointestinal disorder is selected from the group consisting of lymphocytic colitis, microscopic colitis, collagenous colitis, autoimmune enteropathy, allergic gastrointestinal disease and eosinophilic gastrointestinal disease.

11. The method of claim 7, wherein said 6-mercaptapurine metabolite is 6-thioguanine.

12. The method of claim 11, wherein said predetermined toxic level of 6-thioguanine corresponds to a level of about 400 pmol per 8×10^8 red blood cells.

13. The method of claim 11, wherein said toxicity associated with 6-mercaptapurine drug treatment is hematologic toxicity.

14. The method of claim 7, wherein said 6-mercaptapurine metabolite is 6-methyl-mercaptapurine.

15. The method of claim 14, wherein said predetermined toxic level of 6-methyl-mercaptapurine corresponds to a level of about 7000 pmol per 8×10^8 red blood cells.

16. The method of claim 14, wherein said toxicity associated with 6-mercaptapurine treatment is hepatic toxicity.

17. The method of claim 7, wherein said level of 6-mercaptopurine metabolite is determined in red blood cells.

18. The method of claim 17, wherein said level
5 is determined using high pressure liquid chromatography.

19. A method of optimizing therapeutic efficacy and reducing toxicity associated with 6-mercaptopurine drug treatment of an immune-mediated gastrointestinal disorder, comprising:

10 (a) administering a 6-mercaptopurine drug to a subject having said immune-mediated gastrointestinal disorder;

(b) determining a level of 6-thioguanine in said subject having said immune-mediated gastrointestinal
15 disorder; and

(c) determining a level of 6-methyl-mercaptopurine in said subject having said immune-mediated gastrointestinal disorder,

wherein a level of 6-thioguanine less than a
20 predetermined minimal therapeutic level indicates a need to increase the amount of 6-mercaptopurine drug subsequently administered to said subject, thereby increasing therapeutic efficacy,

wherein a level of 6-thioguanine greater than a
25 predetermined toxic level of 6-thioguanine indicates a need to decrease the amount of 6-mercaptopurine drug subsequently administered to said subject, thereby reducing toxicity associated with 6-mercaptopurine drug

treatment of said immune-mediated gastrointestinal disorder, and

wherein a level of 6-methyl-mercaptopurine greater than a predetermined toxic level of

- 5 6-methyl-mercaptopurine indicates a need to decrease the amount of 6-mercaptopurine drug subsequently administered to said subject, thereby reducing toxicity associated with 6-mercaptopurine drug treatment of said immune-mediated gastrointestinal disorder.

- 10 20. The method of claim 19, wherein said immune-mediated gastrointestinal disorder is IBD.

21. The method of claim 20, wherein said subject having IBD is a pediatric subject.

- 15 22. The method of claim 19, wherein said immune-mediated gastrointestinal disorder is selected from the group consisting of lymphocytic colitis, microscopic colitis, collagenous colitis, autoimmune enteropathy, allergic gastrointestinal disease and eosinophilic gastrointestinal disease.

- 20 23. The method of claim 19, wherein said predetermined minimal therapeutic level of 6-thioguanine is a level corresponding to about 230 pmol per 8×10^8 red blood cells.

- 25 24. The method of claim 19, wherein said predetermined toxic level of 6-thioguanine is a level corresponding to about 400 pmol per 8×10^8 red blood cells.

25. The method of claim 19, wherein said predetermined toxic level of 6-methyl-mercaptopurine is a level corresponding to about 7000 pmol per 8×10^8 red blood cells.

5 26. The method of claim 19, wherein said predetermined minimal therapeutic level of 6-thioguanine is a level corresponding to about 230 pmol per 8×10^8 red blood cells, said predetermined toxic level of 6-thioguanine is a level corresponding to about 400 pmol
10 per 8×10^8 red blood cells, and said predetermined toxic level of 6-methyl-mercaptopurine is a level corresponding to about 7000 pmol per 8×10^8 red blood cells.

27. The method of claim 19, wherein said level of 6-thioguanine and said level of
15 6-methyl-mercaptopurine each is determined in red blood cells.

28. The method of claim 27, wherein said level is determined using high pressure liquid chromatography.

29. The method of claim 19, wherein said
20 toxicity associated with 6-mercaptopurine drug treatment is selected from the group consisting of hepatic toxicity and hematologic toxicity.

30. A method of optimizing therapeutic efficacy of 6-mercaptopurine drug treatment of a non-IBD autoimmune disease, comprising:

(a) administering a 6-mercaptopurine drug to a
5 subject having said non-IBD autoimmune disease; and

(b) determining a level of 6-thioguanine in
said subject having said non-IBD autoimmune disease,

wherein a level of 6-thioguanine less than a
minimal therapeutic level indicates a need to increase
10 the amount of 6-mercaptopurine drug subsequently
administered to said subject and

wherein a level of 6-thioguanine greater than a
level corresponding to a predetermined toxic level
indicates a need to decrease the amount of
15 6-mercaptopurine drug subsequently administered to said
subject.

31. The method of claim 30, wherein said
minimal therapeutic level is about 230 pmol per 8×10^8 red
blood cells.

20 32. The method of claim 30, wherein said
predetermined toxic level is about 400 pmol per 8×10^8 red
blood cells.

33. The method of claim 30, wherein said level
of 6-mercaptopurine metabolite is determined in red blood
25 cells.

34. The method of claim 33, wherein said level
is determined using high pressure liquid chromatography.

ABSTRACT OF THE INVENTION

The present invention provides a method of optimizing therapeutic efficacy and reducing toxicity associated with 6-mercaptopurine drug treatment of an
 5 immune-mediated gastrointestinal disorder such as inflammatory bowel disease. The method of the invention includes the step of determining the level of one or more 6-mercaptopurine metabolites in the patient having an immune-mediated gastrointestinal disorder.

10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782
 783
 784
 785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796
 797
 798
 799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000
 1001
 1002
 1003
 1004
 1005
 1006
 1007
 1008
 1009
 1010
 1011
 1012
 1013
 1014
 1015
 1016
 1017
 1018
 1019
 1020
 1021
 1022
 1023
 1024
 1025
 1026
 1027
 1028
 1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048
 1049
 1050
 1051
 1052
 1053
 1054
 1055
 1056
 1057
 1058
 1059
 1060
 1061
 1062
 1063
 1064
 1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073
 1074
 1075
 1076
 1077
 1078
 1079
 1080
 1081
 1082
 1083
 1084
 1085
 1086
 1087
 1088
 1089
 1090
 1091
 1092
 1093
 1094
 1095
 1096
 1097
 1098
 1099
 1100
 1101
 1102
 1103
 1104
 1105
 1106
 1107
 1108
 1109
 1110
 1111
 1112
 1113
 1114
 1115
 1116
 1117
 1118
 1119
 1120
 1121
 1122
 1123
 1124
 1125
 1126
 1127
 1128
 1129
 1130
 1131
 1132
 1133
 1134
 1135
 1136
 1137
 1138
 1139
 1140
 1141
 1142
 1143
 1144
 1145
 1146
 1147
 1148
 1149
 1150
 1151
 1152
 1153
 1154
 1155
 1156
 1157
 1158
 1159
 1160
 1161
 1162
 1163
 1164
 1165
 1166
 1167
 1168
 1169
 1170
 1171
 1172
 1173
 1174
 1175
 1176
 1177
 1178
 1179
 1180
 1181
 1182
 1183
 1184
 1185
 1186
 1187
 1188
 1189
 1190
 1191
 1192
 1193
 1194
 1195
 1196
 1197
 1198
 1199
 1200
 1201
 1202
 1203
 1204
 1205
 1206
 1207
 1208
 1209
 1210
 1211
 1212
 1213
 1214
 1215
 1216
 1217
 1218
 1219
 1220
 1221
 1222
 1223
 1224
 1225
 1226
 1227
 1228
 1229
 1230
 1231
 1232
 1233
 1234
 1235
 1236
 1237
 1238
 1239
 1240
 1241
 1242
 1243
 1244
 1245
 1246
 1247
 1248
 1249
 1250
 1251
 1252
 1253
 1254
 1255
 1256
 1257
 1258
 1259
 1260
 1261
 1262
 1263
 1264
 1265
 1266
 1267
 1268
 1269
 1270
 1271
 1272
 1273
 1274
 1275
 1276
 1277
 1278
 1279
 1280
 1281
 1282
 1283
 1284
 1285
 1286
 1287
 1288
 1289
 1290
 1291
 1292
 1293
 1294
 1295
 1296
 1297
 1298
 1299
 1300
 1301
 1302
 1303
 1304
 1305
 1306
 1307
 1308
 1309
 1310
 1311
 1312
 1313
 1314
 1315
 1316
 1317
 1318
 1319
 1320
 1321
 1322
 1323
 1324
 1325
 1326
 1327
 1328
 1329
 1330
 1331
 1332
 1333
 1334
 1335
 1336
 1337
 1338
 1339
 1340
 1341
 1342
 1343
 1344
 1345
 1346
 1347
 1348
 1349
 1350
 1351
 1352
 1353
 1354
 1355
 1356
 1357
 1358
 1359
 1360
 1361
 1362
 1363
 1364
 1365
 1366
 1367
 1368
 1369
 1370
 1371
 1372
 1373
 1374
 1375
 1376
 1377
 1378
 1379
 1380
 1381
 1382
 1383
 1384
 1385
 1386
 1387
 1388
 1389
 1390
 1391
 1392
 1393
 1394
 1395
 1396
 1397
 1398
 1399
 1400
 1401
 1402
 1403
 1404
 1405
 1406
 1407
 1408
 1409
 1410
 1411
 1412
 1413
 1414
 1415
 1416
 1417
 1418
 1419
 1420
 1421
 1422
 1423
 1424
 1425
 1426
 1427
 1428
 1429
 1430
 1431
 1432
 1433
 1434
 1435
 1436
 1437
 1438
 1439
 1440
 1441
 1442
 1443
 1444
 1445
 1446
 1447
 1448
 1449
 1450
 1451
 1452
 1453
 1454
 1455
 1456
 1457
 1458
 1459
 1460
 1461
 1462
 1463
 1464
 1465
 1466
 1467
 1468
 1469
 1470
 1471
 1472
 1473
 1474
 1475
 1476
 1477
 1478
 1479
 1480
 1481




FIGURE 1






FIGURE 3